

MECHANISMS OF THE REACTION $\pi^- p \rightarrow a_0^0(980)n \rightarrow \pi^0 \eta n$ AT HIGH ENERGIES

N.N. Achasov and G.N. Shestakov

*Laboratory of Theoretical Physics, S.L. Sobolev Institute for Mathematics,
630090, Novosibirsk 90, Russia*

Abstract

Over the last twenty years, it was assumed that the reaction $\pi^- p \rightarrow a_0^0(980)n \rightarrow \pi^0 \eta n$ at high energies is dominated by the b_1 Regge pole exchange. We show that the new Brookhaven and Serpukhov data on the t distributions for this reaction can be explained within the framework of the Regge pole model only if the ρ_2 Regge pole exchange conspiring with its daughter plays a crucial role. The tentative estimates of the absolute $\pi^- p \rightarrow a_0^0(980)n \rightarrow \pi^0 \eta n$ reaction cross section are obtained. The contributions of the one-pion exchange mechanism and the Regge cuts are also estimated.

1. In the $q\bar{q}$ -model (q is a light quark), every rotational excitation with the orbital angular momentum L consists of four nonet: states $^{2S+1}L_J = {}^3L_{L-1}, {}^3L_L$, and ${}^3L_{L+1}$ with charge-parity $C = (-1)^{L+1}$ and state 1L_L with $C = (-1)^L$. But, so far there is a white spot in the lower-lying family with $L = 2$ [1]. The nonstrange members of the 3D_2 nonet with quantum numbers $I^G(J^{PC}) = 1^+(2^{--})$ and $0^-(2^{--})$, i.e. the ρ_2, ω_2 , and ϕ_2 mesons (the masses of which are expected near 1.7 and 1.9 GeV [2,3]), are not yet identified as peaks in corresponding multibody mass spectra [1]. However, the mass distributions are not unique keepers of information on the resonances. The resonance spectrum is also reflected in the Regge behavior of the reaction cross sections at high energies. At present detailed investigations of the reaction $\pi^- p \rightarrow a_0^0(980)n$ at $P_{lab}^{\pi^-} = 18$ and 38 GeV/c are carried out, respectively, at Brookhaven [4] and Serpukhov [5]. This reaction involves only unnatural parity exchanges in the t -channel. This fact is exclusive. Just it permits us to conclude that the observed forward peak in the differential cross section of $\pi^- p \rightarrow a_0^0(980)n$ is dominated by the Regge pole with quantum numbers of the “lost” ρ_2 meson [6].

2. The s -channel helicity amplitudes for the reaction $\pi^-(q_1)p(p_1) \rightarrow a_0^0(980, q_2)n(p_2)$ can be written as

$$M_{\lambda_n \lambda_p} = \bar{u}_{\lambda_n}(p_2) \gamma_5 [-A - \gamma^\mu (q_1 + q_2)_\mu B/2] u_{\lambda_p}(p_1) , \quad (1)$$

where $\lambda_p (= +, -)$ and $\lambda_n (= +, -)$ are the proton and neutron helicities, A and B are the invariant amplitudes depending on $s = (p_1 + q_1)^2$ and $t = (q_1 - q_2)^2$ and free from kinematical singularities. In the c.m. system, $d\sigma/dt = (|M_{++}|^2 + |M_{+-}|^2)/64\pi s |\vec{q}_1|^2$ and, at fixed $t < -1$ GeV² and $s \gg m_N^2$, the helicity amplitudes have the form [6]

$$M_{++} \approx -sB , \quad M_{+-} \approx \sqrt{t_{min} - t} A . \quad (2)$$

As it is well known, the t -channel helicity amplitudes with definite parity and free from kinematical singularities are most suitable for Reggeization [7]. These amplitudes for the

reaction $\pi^- a_0^0 \rightarrow \bar{p}n$ have the form

$$G_{++} = tA + m_N(m_{a_0}^2 - m_\pi^2)B, \quad G_{+-} = B. \quad (3)$$

In the Regge region, G_{++} and G_{+-} behave like s^α and $s^{\alpha-1}$ [6]. It is easy to show that the amplitude G_{++} has to contain the Regge pole exchange with $I = 1$, $G = +1$, signature $\tau = -1$, and “naturalness” $\tau P = -1$. The high-lying Regge trajectory with such quantum numbers is the b_1 trajectory (the well-known $b_1(1235)$ meson is its lower-lying representative). The second independent amplitude G_{+-} has to contain the Regge pole exchanges with $I = 1$, $G = +1$, $\tau = +1$, and $\tau P = -1$ and here the ρ_2 Regge trajectory is a leading one. In the physical region of the s -channel, the b_1 and ρ_2 Regge pole contributions can be written as $G_{++}^{b_1} = \beta_{b_1}(t)(s/s_0)^{\alpha_{b_1}(t)} i e^{-i\pi\alpha_{b_1}(t)/2}$ and $G_{+-}^{\rho_2} = \beta_{\rho_2}(t)(s/s_0)^{\alpha_{\rho_2}(t)-1} e^{-i\pi\alpha_{\rho_2}(t)/2}$, where $\beta(t)$, $\alpha(t)$, and complex factors are residues, trajectories, and signature factors of the corresponding Regge poles, and $s_0 = 1 \text{ GeV}^2$. As it is now evident from Eqs. (2) and (3), the leading contributions to the s -channel helicity amplitudes M_{++} and M_{+-} at high energies are caused by the ρ_2 and b_1 Regge exchanges, respectively.

Expressing the invariant amplitudes A and B in terms of G_{++} and G_{+-} ,

$$A = [G_{++} - m_N(m_{a_0}^2 - m_\pi^2)G_{+-}]/t, \quad B = G_{+-}, \quad (4)$$

we find the $1/t$ singular behavior for the invariant amplitude A . To avoid this singularity, one has to impose a further constraint equation between the t -channel helicity amplitudes, namely: at $t = 0$, $G_{++} = m_N(m_{a_0}^2 - m_\pi^2)G_{+-}$. It is of interest to satisfy the analyticity of A at $t = 0$ assuming the different Regge pole models.

If we have the b_1 and ρ_2 exchanges only and assume that the amplitudes $G_{++}^{b_1}$ and $G_{+-}^{\rho_2}$ do not vanish as $t \rightarrow 0$, we obtain two relations, $\alpha_{b_1}(0) = \alpha_{\rho_2}(0) - 1$, and $\beta_{b_1}(0) = m_N(m_{a_0}^2 - m_\pi^2)\beta_{\rho_2}(0)$, the first of which is rather silly because, at the usual values of $\alpha_{b_1}(0) \approx -(0.05 \div 0.3)$, it requires $\alpha_{\rho_2}(0) \approx 0.95 \div 0.7$. For the ρ_2 trajectory heaving unnatural parity, this is evidently ruled out. Of course, in order for the amplitude A to be regular for $t \rightarrow 0$, one can accept that the amplitudes $G_{++}^{b_1}$ and $G_{+-}^{\rho_2}$ are separately proportional to t . But, in this case, a dip in $d\sigma/dt$ in the forward direction is predicted by Eqs. (2) and (3). On the contrary, the Brookhaven and Serpukhov experiments [4,5] show a clear forward peak. This means that the amplitude M_{++} with quantum numbers of the ρ_2 exchange does not vanish as $t \rightarrow 0$. In the framework of the Regge pole model, this can be attended only in the case of a conspiracy of the ρ_2 Regge trajectory with its daughter one (d), which has to have quantum numbers of the b_1 exchange. Let us write down the contribution of such a daughter trajectory near $t = 0$ in the form $G_{++}^d = \beta_d(t)(s/s_0)^{\alpha_d(t)} i e^{-i\pi\alpha_d(t)/2}$. Then, the amplitude A should be regular at $t = 0$ if the following relations for the ρ_2 , d , and b_1 exchanges are valid: $\alpha_d(0) = \alpha_{\rho_2}(0) - 1$, $\beta_d(0) = m_N(m_{a_0}^2 - m_\pi^2)\beta_{\rho_2}(0)$, $\beta_{\rho_2}(0) \neq 0$, and $\beta_{b_1}(t) \sim t$. Thereby the amplitude M_{++} dominated by the ρ_2 exchange does not vanish as $t \rightarrow 0$ and can lead to the observed forward peak. Note that asymptotically (at large s) the daughter contribution and nonasymptotic contribution of the ρ_2 trajectory (which behave as $\sim s^{\alpha_{\rho_2}-1}$) to the amplitude A and consequently to the amplitude M_{+-} can all be neglected.

Thus, in the model with the b_1 and conspiring ρ_2 Regge poles, the data on the t distributions at fixed s can be fitted to the form

$$dN/dt = C_1 e^{\Lambda_1 t} + (t_{min} - t) C_2 e^{\Lambda_2 t}, \quad (5)$$

where the first term corresponds to the contribution of the amplitude M_{++} with the conspiring ρ_2 exchange and the second term corresponds to the contribution of the amplitude M_{+-} with the b_1 exchange. The Brookhaven data for $-t_{min} < -t < 0.6 \text{ GeV}^2$ [4] are fitted by Eq. (5) equally well both with and without the b_1 exchange. The fit with $C_2 \equiv 0$ in Eq. (5) gives $C_1 = 129 \text{ events/GeV}^2$, $\Lambda_1 = 4.7 \text{ GeV}^{-2}$, and $\chi^2/\text{n.d.f.} \approx 15.9/22$. It is shown in Fig. 1 by the solid curve. The fit to these data to the full form in Eq. (5) gives $C_1 = 131 \text{ events/GeV}^2$, $\Lambda_1 = 7.6 \text{ GeV}^{-2}$, $C_2 = 340 \text{ events/GeV}^4$, $\Lambda_2 = 5.8 \text{ GeV}^{-2}$, and $\chi^2/\text{n.d.f.} \approx 15.9/20$. The corresponding curve is practically the same as the solid one in Fig. 1. The dashed and dotted curves in Fig. 1 show the ρ_2 and b_1 contributions separately. The latter yields approximately 34% of the integrated cross section.

In order to have an idea of the absolute value of the $\pi^-p \rightarrow a_0^0(980)n \rightarrow \pi^0\eta n$ cross section, it has been estimated by us in Ref. [6] at $P_{lab}^{\pi^-} = 18 \text{ GeV}/c$ using the available data on the reaction $\pi^-p \rightarrow a_2^0(1320)n$ and the Brookhaven data [4] on the $\pi^0\eta$ mass spectrum in $\pi^-p \rightarrow \pi^0\eta n$. Our tentative estimate is: $\sigma(\pi^-p \rightarrow a_0^0(980)n \rightarrow \pi^0\eta n) \approx 200 \text{ nb}$ and $[d\sigma/dt(\pi^-p \rightarrow a_0^0(980)n \rightarrow \pi^0\eta n)]_{t \approx 0} \approx 940 \text{ nb/GeV}^2$. Note that (6–15)% of the integrated cross section can be caused by the one-pion exchange mechanism which arises owing to the $f_0(980) - a_0^0(980)$ mixing violating isotopic invariance [6]. Also the $a_2\pi$, ρb_1 , ωa_1 , $\rho\rho P$, and $a_2 a_2 P$ Regge cut contributions (where P is the Pomeron) to the amplitude M_{++} have been estimated in Ref. [6] and the conclusion has been made that they are small.

Note that the early estimate of the $a_0(980)$ production [8] assumed the b_1 exchange dominance and consequently a dip in $d\sigma/dt$ in the forward direction.

3. Let us note two practical experimental problems. *i)* It is important to determine rather accurately the parameters of the simplest Regge pole model. To do so it is needed the good data on $d\sigma/dt(\pi^-p \rightarrow a_0^0(980)n)$ at several appreciably different energies. First of all, we have in mind the energies of the π^- beams at Serpukhov, Brookhaven, and KEK. *ii)* It would be very interesting to find some signs of the ρ_2 state and its daughter state having the b_1 meson quantum numbers, for example, in the $a_0\pi$, $\omega\pi$, and $A_2\pi$ mass spectra around 1.7 GeV.

References

- [1] Barnett, R.M. et al. (Particle Data Group), *Phys. Rev. D* **54**, 46, 99 (1996).
- [2] Burkhardt, H., and Hey, A.J.G., *Nucl. Phys. B* **108**, 514 (1976).
- [3] Kokoski, R., and Isgur, N., *Phys. Rev. D* **35**, 907 (1987).
- [4] Dzierba, R.A., in *Proceedings of the Second Workshop on Physics and Detectors for DAΦNE'95*, Frascati, 1995, *Frascati Physics Series* **4**, 99 (1996).
- [5] Sadovsky, S.A., Talk presented at this Conference.
- [6] Achasov, N.N., and Shestakov, G.N., *Phys. Rev. D* **56**, 212 (1997).
- [7] Bertocchi, L., in *Proceedings of the Heidelberg International Conference on Elementary Particles*, Heidelberg, 1967, North-Holland Publ. Co., 1968, p. 197.
- [8] Irving, A.C., *Phys. Lett. B* **70**, 217 (1977).

Figure caption

Fig. 1. The t distribution for the reaction $\pi^- p \rightarrow a_0^0(980)n \rightarrow \pi^0 \eta n \rightarrow 4\gamma n$ at 18 GeV/c measured at Brookhaven [4]. The fits are described in the text.




Fig. 1.